

Home Search Collections Journals About Contact us My IOPscience

Discretized rotation has infinitely many periodic orbits

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2013 Nonlinearity 26 871

(http://iopscience.iop.org/0951-7715/26/3/871)

View the table of contents for this issue, or go to the journal homepage for more

Download details: IP Address: 193.6.168.18 The article was downloaded on 14/02/2013 at 12:47

Please note that terms and conditions apply.

Nonlinearity 26 (2013) 871-880

Discretized rotation has infinitely many periodic orbits

Shigeki Akiyama¹ and Attila Pethő²

 ¹ Institute of Mathematics, University of Tsukuba, Tennodai 1-1-1, Tsukuba, Ibaraki, 305-8571 Japan
 ² Department of Computer Science, University of Debrecen, PO Box 12, H-4010 Debrecen,

² Department of Computer Science, University of Debrecen, PO Box 12, H-4010 Debrecen, Hungary

E-mail: akiyama@math.tsukuba.ac.jp and petho.attila@inf.unideb.hu

Received 15 August 2012, in final form 11 December 2012 Published 13 February 2013 Online at stacks.iop.org/Non/26/871

Recommended by C Liverani

Abstract

For a fixed $\lambda \in (-2, 2)$, we study a family of discretized rotation on \mathbb{Z}^2 defined by

 $(x, y) \mapsto (y, -\lfloor x + \lambda y \rfloor).$

We prove that this reversible dynamics has infinitely many periodic orbits.

Mathematics Subject Classification: 37E99, 37A45, 37P35

(Some figures may appear in colour only in the online journal)

1. Introduction

Space discretization of dynamical systems has attracted considerable interest from researchers [6, 9, 18, 24, 27]. One motivation is to understand the distance between the original dynamics and its computer simulation through a discretized model. In this paper, we are interested in discretized planar rotation. It is a very simple system, but yet we know surprisingly little about this discretized system. We start with a conjecture studied by many authors, for example, in [11, 20, 25] and from the point of view of a shift radix system in [1].

Conjecture. For all fixed $-2 < \lambda < 2$, all integer sequences (a_n) defined by

$$0 \leqslant a_{n+2} + \lambda a_{n+1} + a_n < 1 \tag{1}$$

with the initial value $(a_0, a_1) \in \mathbb{Z}^2$ are periodic.

For $(x, y) = (a_n, a_{n+1})$, we have

$$(a_{n+1}, a_{n+2}) = (y, -|x + \lambda y|),$$

0951-7715/13/030871+10\$33.00 © 2013 IOP Publishing Ltd & London Mathematical Society Printed in the UK & the USA 871

and it defines a map $F : (x, y) \mapsto (y, -\lfloor x + \lambda y \rfloor)$ on \mathbb{Z}^2 . In other words, we are interested in the dynamics F on \mathbb{Z}^2 :

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 0 & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ \langle \lambda y \rangle \end{pmatrix},$$
(2)

where $\langle x \rangle = x - \lfloor x \rfloor$. Since the eigenvalues of the matrix are two conjugate complex numbers of modulus one, this dynamics can be regarded as a rotation with invariant confocal ellipses, acting on the lattice \mathbb{Z}^2 . After the rotation of angle θ with $\lambda = -2\cos\theta$, we translate by a small vector to make the image lie in \mathbb{Z}^2 . An affine equivalent formulation using Euclidean rotation is found in the next section. From the shape of the inequality in (1), the dynamics (2) is reversible, i.e. F(x, y) = (y, z) implies F(z, y) = (y, x). Thus we have $\phi F^{-1} = F\phi$ with $\phi(x, y) = (y, x)$ and F is a bijection on \mathbb{Z}^2 . In other words, F is a composition of the two involutions $F\phi$ and ϕ .

The conjecture is supported by numerical experiments [1, 24]. It is also expected from a heuristic perspective: the cumulation of errors of F^n from the exact $n\theta$ rotation is expected to be small and it is seemingly impossible to avoid hitting the same lattice points. The cumulative error bound is discussed in [18, 26]. However this problem is notorious, and our knowledge is limited. We only know that conjecture 1 holds for 11 values $\lambda = 0, \pm 1, (\pm 1 \pm \sqrt{5})/2, \pm \sqrt{2}, \pm \sqrt{3}, \text{ see } [1, 2, 20].$ Apart from three trivial cases $0, \pm 1, 1, 1, 2, 20$ the proof is highly non-trivial and uses the self-inducing structure found in the associated planar piecewise isometry when θ/π is rational and λ is quadratic. If θ/π is rational, then we can embed the problem into piecewise isometry acting on a certain higher dimensional torus (see [17, 20], and also [4, 5] for a connection to digital filters). Piecewise isometries have zero entropy [12], but we know little about their periodic orbits [14]. It is noteworthy that a certain piecewise isometry generated by 7-fold rotation in the plane is governed by several self-inducing structures [3, 15, 19], but it is irrelevant to the map F. If λ is a rational number whose denominator is the power of a prime, then the dynamics is understood as the composition of p-adic rotation and symbolic shift in [10], but it seems difficult to extract information on periodic orbits through this embedding. At this stage, we are interested in giving a non-trivial general statement for this dynamics. In this paper, we will show

Theorem 1. For all fixed $\lambda \in (-2, 2)$ there are infinitely many periodic orbits of the dynamics (2) on \mathbb{Z}^2 .

More precisely, we prove that there are infinitely many *symmetric* periodic orbits (see section 3 for the definition). See corollary 3 in section 6, for a qualitative statement. Theorem 1 is new for all λ except the above 11 values, and gives another support of the conjecture. Note that the idea of observing symmetric periodic orbits dates back to Birkhoff, who showed the existence of infinitely many symmetric periodic orbits for the restricted three body problem [7, 8], whose dynamics is composed of two involutions.

Adding a counting technique of lattice points in number theory, we can generalize theorem 1 to the sequences generated by

$$-\eta \leqslant a_{n+2} + \lambda a_{n+1} + a_n < 1 - \eta$$

for a fixed $\eta \in \mathbb{R}$, as in section 6, theorem 2. It covers a significant class of discretized rotations, but we do not know how large this two parameter family is within the set of all invertible discretized rotations in \mathbb{Z}^2 , up to conjugacy.

We say that p = p(x, y) > 0 is the *period* of $(F^n(x, y))_{n \in \mathbb{Z}}$, if it is the smallest positive integer p with $F^p(x, y) = (x, y)$. If there is no such p, then p(x, y) is not defined. It is remarkable that the distribution of periodic orbits drastically changes if θ/π is irrational or rational. We have:

Lemma 1. Let θ/π be irrational and p be a positive integer. Then there are only finitely many periodic orbits of period p.

This fact follows from theorem 2.1 of [24] but we give a quick proof in section 5. On the other hand, if θ/π is rational, in view of the above torus embedding, it is natural to obtain infinitely many periodic orbits of period p, which fall into the same period cell. Theorem 4.3 in [1] gives a concrete example of infinite periodic orbits of period p for $\theta = (1 - 1/p)\pi$ with an odd prime p.

By theorem 1 and lemma 1, we know that there exist arbitrary large periods if θ/π is irrational. We expect the same to hold for all $\lambda \neq 0, \pm 1$, but there are proofs only for the above eight quadratic cases.

2. Setting and strategy

Let $\lambda = -2\cos\theta$ where θ is a real number in $(0, \pi)$ and let $Q = \begin{pmatrix} -\sin\theta & \cos\theta\\ 0 & 1 \end{pmatrix}$. Our

transformation on \mathbb{Z}^2 : $(x, y) \mapsto (X, Y)$ is written as

$$\begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & -\lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ \mu \end{pmatrix}$$

b. Since

with $\mu \in [0, 1)$. Since

$$Q\begin{pmatrix}\cos\theta & -\sin\theta\\\sin\theta & \cos\theta\end{pmatrix}Q^{-1} = \begin{pmatrix}0 & 1\\-1 & -\lambda\end{pmatrix},$$

we view this algorithm as

$$Q^{-1}\begin{pmatrix} X\\ Y \end{pmatrix} = \begin{pmatrix} \cos\theta & -\sin\theta\\ \sin\theta & \cos\theta \end{pmatrix} Q^{-1}\begin{pmatrix} x\\ y \end{pmatrix} + Q^{-1}\begin{pmatrix} 0\\ \mu \end{pmatrix}.$$
 (3)

Thus, it is the dynamics acting on the lattice $\mathcal{L} = \begin{pmatrix} -\csc\theta\\ 0 \end{pmatrix} \mathbb{Z} + \begin{pmatrix} \cot\theta\\ 1 \end{pmatrix} \mathbb{Z}$ written as the composition of the Euclidean rotation of angle θ followed by a small translation

$$v \mapsto v + \mu \begin{pmatrix} \cot \theta \\ 1 \end{pmatrix}$$

with $\mu \in [0, 1)$. Let *R* be a positive real number and B(R) be a ball of radius *R* centred at the origin. Define a *trap region* T(R) by

$$T(R) = \left\{ x + y \begin{pmatrix} \cot \theta \\ 1 \end{pmatrix} \middle| x \in B(R), y \in [0, 1) \right\} \setminus B(R).$$

The situation is demonstrated in figure 1.

Now we explain the strategy of the proof. It is clear from the description of the dynamics that every unbounded orbit starting from a point in $\mathcal{L} \cap B(R)$ must visit at least once the trap region T(R). Assume that there are only finitely many periodic orbits of (3). Since we are dealing with dynamics on the lattice \mathcal{L} , periodicity of an orbit is equivalent to its boundedness. We argue by contradiction: the assumption that there are just a finite number of periodic orbits would imply that, for $R \gg 1$, the number of lattice points in T(R) are strictly less than the number of symmetric unbounded orbits starting from B(R), which is impossible.

3. Lower bound of unbounded orbits

Symmetric periodic orbits of time-reversal dynamics have been studied for a long time. We shall make use of a well-known property. Let (a_n) be a bi-infinite integer sequence and b be



Figure 1. Trap region.

an integer. We say that (a_n) is periodic, if $a_{n+b} = a_n$ holds for all n. The sequence (a_n) is symmetric at b/2, if $a_{b-n} = a_n$ holds for all n, and it is *doubly symmetric* if it is symmetric at $b_1/2$ and $b_2/2$ with $b_1 \neq b_2$.

Lemma 2. If a sequence is doubly symmetric then it is periodic. Moreover symmetric and periodic sequences are doubly symmetric.

Proof. Assume that (a_n) is symmetric at b_1 and b_2 with $b_1 \neq b_2$. Then $a_{n+b_2-b_1} = a_{b_1-n} = a_n$. Let (a_n) be periodic of period *b* and symmetric at *c*. Then $a_{c-b-n} = a_{n+b} = a_n$.

In fact, this lemma is a restatement of theorems 1 and 2 in [13], where symmetric periodic orbits of time reversal dynamics composed of two involutions are studied. The reader can find precise description on such periodic orbits there.

Let $(x, y) \in \mathbb{Z}^2$. To the bi-infinite orbit $(F^n(x, y))_{n \in \mathbb{Z}}$ we can associate uniquely the bi-infinite sequence (a_n) consisting of the 1st coordinates of the elements of the orbit. It is clear that $(F^n(x, y))$ is periodic if and only if (a_n) is periodic. Hereafter we identify the orbit $(F^n(x, y))$ and the bi-infinite sequence (a_n) and say that an orbit $(F^n(x, y))$ is symmetric if (a_n) is so. Assume that (a_n) is symmetric at b/2. If b is odd, then $a_{(b-1)/2} = a_{(b+1)/2}$ and the orbit is of the form:

$$\ldots, c_3, c_2, c_1, X, X, c_1, c_2, c_3, \ldots$$

with some $X \in \mathbb{Z}$ and a sequence $(c_n) \subset \mathbb{Z}$. Clearly (c_n) is determined by X. Such orbits are in Fix (ϕ) , the set of orbits fixed by the involution ϕ . If b is even, then $a_{b/2-1} = a_{b/2+1}$ and the orbit is of the form

 $\ldots, c_3, c_2, c_1, X, Y, X, c_1, c_2, c_3, \ldots$

for some $X, Y \in \mathbb{Z}$ and $(c_n) \subset \mathbb{Z}$. Of course (c_n) is determined by X and Y. These orbits belong to Fix $(F\phi)$, the set fixed by the other involution $F\phi$. By lemma 2, each periodic orbit of $(F^n(x, y))$ is doubly symmetric. It belongs to the same fix point set if the period is even, and to a different fix point sets if the period is odd.

Remark 1. Not all orbits are symmetric. For example, if $\lambda = (1 + \sqrt{5})/2$ then we have

$$(-1,4) \rightarrow (4,-6) \rightarrow (-6,5) \rightarrow (5,-3) \rightarrow (-3,-1) \rightarrow (-1,4).$$

We do not know a way to estimate the number of asymmetric orbits from below.

By our assumption of *reductio ad absurdum*, there exist only a finite number, say C_1 , of periodic orbits. For any R, the number of points in $\mathcal{L} \cap B(R)$ whose orbits are periodic is less than C_1 .

3.1. Unbounded orbits in $Fix(\phi)$

Let (a_n) be an unbounded orbit in Fix (ϕ) . By lemma 2, (a_n) cannot be doubly symmetric, which implies that two orbits starting from different fixed points in \mathcal{L} never intersect. Apart from a finite number of exceptions, points of the form

$$X\begin{pmatrix} -\csc\theta\\ 0 \end{pmatrix} + X\begin{pmatrix} \cot\theta\\ 1 \end{pmatrix} \in \mathcal{L} \cap B(R)$$

generate distinct unbounded orbits. From

$$X^2 \left(-\csc\theta + \cot\theta\right)^2 + X^2 \leqslant R^2,$$

we conclude that there are at least $2R \cos(\theta/2) - C_1$ unbounded orbits in Fix(ϕ) starting from $\mathcal{L} \cap B(R)$.

3.2. Unbounded orbits in $Fix(F\phi)$

Similarly, by lemma 2, (a_n) cannot be doubly symmetric. So, by (1), our task is to count the number of the pairs (X, Y) which satisfy

$$0 \leqslant X + \lambda Y + X < 1 \tag{4}$$

and

$$X\begin{pmatrix} -\csc\theta\\ 0 \end{pmatrix} + Y\begin{pmatrix} \cot\theta\\ 1 \end{pmatrix} \in B(R).$$
(5)

For this computation, we substitute the inequality (4) by

$$-1 \leqslant X + \lambda Y + X < 1 \tag{6}$$

and count the number of pairs (X, Y) satisfying (6) and (5). It is clear that for a fixed Y, there is a unique X which satisfies (6). Since $X = Y \cos \theta + \varepsilon$ with $|\varepsilon| \le 1/2$, we have

$$(Y\cot\theta - X\csc\theta)^2 + Y^2 = Y^2 + \frac{\varepsilon^2}{\sin^2\theta} \leqslant R^2$$
(7)

from (5), and we have at least $2R - C_2$ such points with a non-negative constant C_2 .

If
$$\lambda$$
 is irrational, then there is no $(X, Y) \neq (0, 0)$ which satisfies either

$$-1 = 2X + \lambda Y$$
 or $0 = 2X + \lambda Y$.

Using the symmetry $(X, Y) \leftrightarrow (-X, -Y)$, we see that the number of (X, Y) with (5) and

$$-1 \leq X + \lambda Y + X < 0$$

is exactly one less than the number of (X, Y) with (4) and (5), which counts the origin. Thus the number of (X, Y) having (4) and (5) is at least $R - C_2/2$.

If λ is rational, then we additionally have to take care of the points (X, Y) on the line $-1 = 2X + \lambda Y$ and $0 = 2X + \lambda Y$. However we can easily show that the number of (X, Y) with (5) on the line $-1 = 2X + \lambda Y$ and the one on the line $0 = 2X + \lambda Y$ differ only by some constant. Thus in any case, there are at least $R - C_3$ unbounded orbits in Fix $(F\phi)$ starting from $\mathcal{L} \cap B(R)$ with a non-negative constant C_3 .

4. Lattice points in the trap region

By construction, if the trap region T(R) and the line $x = y \cot \theta + c$ have a non-empty intersection, then it is a half-open interval of length $\csc \theta$. Thus if the line $x = y \cot \theta + c$ intersects $\mathcal{L} \cap T(R)$ then it is a single point. The lattice \mathcal{L} is covered by a family of parallel lines:

$$\Xi = \{ x = y \cot \theta - k \csc \theta \mid k \in \mathbb{Z} \}.$$

We easily see that the distance between adjacent lines of Ξ is 1. Thus there are exactly $2\lfloor R \rfloor + 1$ points in $\mathcal{L} \cap T(R)$.

Remark 2. If $\theta < 2\pi/3$, then $2R \cos(\theta/2) - C_1 + R - C_3 > 2R + 1$ holds for sufficiently large *R* and we immediately obtain the desired contradiction.

Let us take into account the reversibility of *F*. Since we are dealing with unbounded symmetric orbits starting from $\mathcal{L} \cap B(R)$, if $(a_n, a_{n+1}) = (C, D)$ then there is an index *m* such that $(a_m, a_{m+1}) = (D, C)$. Let $\Phi : \mathcal{L} \mapsto \mathcal{L}$ be defined as follows:

$$\Phi: x\begin{pmatrix} -\csc\theta\\ 0 \end{pmatrix} + y\begin{pmatrix} \cot\theta\\ 1 \end{pmatrix} \mapsto y\begin{pmatrix} -\csc\theta\\ 0 \end{pmatrix} + x\begin{pmatrix} \cot\theta\\ 1 \end{pmatrix}.$$

If an orbit visits $\Phi(T(R)) \cap T(R)$ then the number of visits is at least two. In other words, we only have to count the number of lattice points up to this symmetry by Φ in T(R).

The mapping Φ is the reflection with respect to the vector $\begin{pmatrix} -\csc\theta + \cot\theta\\ 1 \end{pmatrix}$, because the

two vectors $\begin{pmatrix} -\csc\theta\\ 0 \end{pmatrix}$ and $\begin{pmatrix} \cot\theta\\ 1 \end{pmatrix}$ have the same length. Thus the reflection Φ leaves the vector $\begin{pmatrix} -\sin(\theta/2)\\ \cos(\theta/2) \end{pmatrix} = \begin{pmatrix} -\csc\theta + \cot\theta\\ 1 \end{pmatrix} \cos(\theta/2)$ invariant.

To make computation easy, we rotate T(R) and \mathcal{L} by $-\theta$ and present the situation in figure 2. T(R)', \mathcal{L}' are the images by this rotation and Ψ is the corresponding reflection. Then every line y = k with $k \in \mathbb{Z} \cap [-R, R]$ contains a single point in $\mathcal{L}' \cap T(R)'$ and the reflection Ψ leaves the vector

$$\begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} -\sin(\theta/2) \\ \cos(\theta/2) \end{pmatrix} = \begin{pmatrix} \sin(\theta/2) \\ \cos(\theta/2) \end{pmatrix}$$

invariant.

So our task is to estimate from above the number of lattice points in $\mathcal{L}' \cap T(R)'$ which are below the line $y = x \cot(\theta/2)$. The intersection of the line and the boundary of T(R)' with the largest y-coordinate is

$$\left(\sqrt{R^2\sin^2\left(\frac{\theta}{2}\right) - \frac{1}{4}} + \frac{1}{2}\tan\frac{\theta}{2}, \cot\frac{\theta}{2}\sqrt{R^2\sin^2\left(\frac{\theta}{2}\right) - \frac{1}{4}} + \frac{1}{2}\right)$$



Figure 2. Symmetry of the trap region.

and we have

$$\cot\frac{\theta}{2}\sqrt{R^2\sin^2\left(\frac{\theta}{2}\right)-\frac{1}{4}}+\frac{1}{2}=R\cos\left(\frac{\theta}{2}\right)+\frac{1}{2}+O\left(\frac{1}{R}\right).$$

We count the number of points whose *y*-coordinate does not exceed this value, i.e. the points in the shaded part in figure 2. Thus the number of lattice points up to symmetry in $\mathcal{L} \cap T(R)$ is bounded from above by $R + R \cos(\theta/2) + C_4$ with a non-negative constant C_4 .

5. Proof of theorem 1 and lemma 1

From the assumption that there are only finitely many periodic orbits, we derived several estimates in the previous sections. By lemma 2, unbounded orbits in $Fix(\phi)$ and those in $Fix(F\phi)$ have no intersection. Thus

$$2R\cos(\theta/2) - C_1 + R - C_3$$

distinct unbounded orbits must visit T(R) and there are only

$$R + R\cos(\theta/2) + C_4$$

lattice points in $\mathcal{L} \cap T(R)$ up to symmetry. However

$$2R\cos(\theta/2) - C_1 + R - C_3 \leqslant R + R\cos(\theta/2) + C_4$$

does not hold for sufficiently large R. The proof of theorem 1 is finished.

Let us show lemma 1. Assume that there are infinitely many (x, y) so that $F^p(x, y) = (x, y)$ with p > 2. By induction using (3), we have

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} \cos p\theta & -\sin p\theta \\ \sin p\theta & \cos p\theta \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \sum_{i=1}^{p} v_i$$

where $\begin{pmatrix} u \\ v \end{pmatrix} \in \mathcal{L}$ and $||v_i|| \leq \csc \theta$. Here $||\cdot||$ is the Euclidean norm. However if $\begin{pmatrix} u \\ v \end{pmatrix}$ is sufficiently large, then

$$\begin{pmatrix} u \\ v \end{pmatrix} - \begin{pmatrix} \cos p\theta & -\sin p\theta \\ \sin p\theta & \cos p\theta \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} \| \ge c \| \begin{pmatrix} u \\ v \end{pmatrix} \| > p \csc \theta.$$

Here c is computed as the operator norm:

$$c = \frac{1}{\left\| \begin{pmatrix} 1 - \cos p\theta & \sin p\theta \\ -\sin p\theta & 1 - \cos p\theta \end{pmatrix}^{-1} \right\|} = 2 \left| \sin \left(\frac{p\theta}{2} \right) \right|$$

which is positive since θ/π is irrational. This gives a contradiction.

6. Generalization

One can generalize the result to the sequences defined by:

 $-\eta \leqslant a_{n+2} + \lambda a_{n+1} + a_n < 1 - \eta,$

with $\eta \in \mathbb{R}$. These kind of interval shifts are studied, for example, in [17, 21]. In complete analogy to our main result, we have

Theorem 2. For a fixed $\lambda \in (-2, 2)$ and $\eta \in \mathbb{R}$, there are infinitely many periodic orbits of *the dynamics*

$$(x, y) \to (y, -\lfloor \lambda y + x + \eta \rfloor)$$
(8)

on \mathbb{Z}^2 .

We can deduce a qualitative statement:

Corollary 3. There is a positive constant C depending on λ and η such that within B(R), there are at least CR periodic orbits of (8).

Hereafter we sketch the proof of theorem 2 and corollary 3. Writing $\kappa = \eta/(2 + \lambda)$, the inequality becomes

$$0 \leq (a_{n+2} + \kappa) + \lambda(a_{n+1} + \kappa) + (a_n + \kappa) < 1$$

Therefore by substituting \mathcal{L} with $\mathcal{L}' = \mathcal{L} + Q^{-1} \binom{\kappa}{\kappa}$, our algorithm has exactly the same shape as (3). Although the error term becomes worse than the one in section 4, we can show that the number of lattice points of \mathcal{L}' up to symmetry within the trap region is

 $R + R\cos(\theta/2) + O(R^{2/3+\epsilon})$

for any positive constant ϵ . Here we used the method of Vinogradov to count the number of lattice points in the cylindrical region bounded by curves of positive curvature, for example, see [23, pp 8–22] or [16, 22].

Similarly to section 3, there are $2R \cos(\theta/2) - C_1$ unbounded orbits in Fix(ϕ). We count unbounded orbits in Fix($F\phi$), i.e. the number of (X, Y), which satisfy

$$\lambda Y/2 \mod 1 \cap [-\eta/2, (1-\eta)/2) \neq \emptyset \tag{9}$$

and (7).

If λ is irrational, then $(\lambda/2)Y \mod 1$ is uniformly distributed and the number of such Y is R + o(R). When λ is rational, put $\lambda/2 = p/q$ with (p, q) = 1. Then $(\lambda/2)Y \equiv i/q \mod 1$ for $i \in \{0, 1, \dots, q-1\}$ with the same frequency 1/q.

Let us study the case that q is even. Since

 $\{i/q \mod 1\} \cap [-\eta/2, (1-\eta)/2)$

has cardinality q/2, the number of points with (9) and (7) is again R + o(R). Once we have this estimate R + o(R) then

$$2R\cos(\theta/2) - C_1 + R + o(R) \leqslant R + R\cos(\theta/2) + o(R^{2/3+\epsilon})$$

does not hold for sufficiently large *R*. Here C_1 appears only once in the left side, because it is the number of periodic orbits of the system. We obtain the contradiction for $R \gg 1$.

It remains to show the case when q is odd. Then

$$\{i/q \mod 1\} \cap [-\eta/2, (1-\eta)/2)$$

has cardinality either (q - 1)/2 or (q + 1)/2 depending on η . Thus the number of unbounded orbits in Fix $(F\phi)$ is bounded from below by R - R/q + o(R). Thus we have to show that

$$2R\cos(\theta/2) - C_1 + R - R/q + o(R) > R + R\cos(\theta/2) + o(R^{2/3+\epsilon})$$

for large R. This is valid because

$$\cos(\theta/2) > 1/q$$

holds for q > 2, since $\cos(\theta/2) = \sqrt{(1 + \cos(\theta))/2} = \sqrt{(1 - p/q)/2}$.

The validity of the statement of corollary 3 is invariant under affine transformations fixing the origin, up to appropriate changes the constant C. To prove corollary 3, note that we may take

$$C_1 = (1 - \varepsilon)R\left(\cos(\theta/2) - 1/q\right)$$

with a small $\varepsilon > 0$ in the above proof to get the same contradiction. Here the term -1/q is necessary only in the last case of the proof.

Acknowledgments

We are grateful to Franco Vivaldi for giving us comments and references for the earlier version of this article. The first author moved from Niigata University to their current address in August 2012. The paper was written when the second author was visiting Niigata University as a long term research fellow of JSPS. Both authors wish to express their gratitude to all the staff at the Department of Mathematics, Niigata University and the support from JSPS.

The authors are supported by the Japanese Society for the Promotion of Science (JSPS), Grant in aid 21540010 and Invitation Fellowship Program FY2011, L-11514.

References

- Akiyama S, Brunotte H, Pethő A and Steiner W 2006 Remarks on a conjecture on certain integer sequences *Periodica Math. Hungarica* 52 1–17
- [2] Akiyama S, Brunotte H, Pethő A and Steiner W 2008 Periodicity of certain piecewise affine planar maps *Tsukuba J. Math.* 32 1–55
- [3] Akiyama S and Harriss E 2013 Pentagonal domain exchange Discrete Contin. Dyn. Syst. at press
- [4] Ashwin P 1997 Elliptic behaviour in the sawtooth standard map *Phys. Lett.* A 232 409–16
- [5] Ashwin P, Chambers W and Petrov G 1997 Lossless digital filters overflow oscillations: approximations of invariant fractals Int. J. Bifurcation Chaos 7 2603–10
- [6] Berthé V and Nouvel B 2007 Discrete rotations and symbolic dynamics Theor. Comput. Sci. 380 276-85
- [7] Birkhoff G D 1915 The restricted problem of three bodies Rendiconti Circolo Matematico Palermo 39 265-334
- [8] Birkhoff G D 1966 Dynamical Systems vol 9 (American Mathematical Society Colloquium Publications) (Providence, RI: American Mathematical Society)

- [9] Blank M 1994 Pathologies generated by round-off in dynamical systems Physica D 78 93-114
- [10] Bosio D and Vivaldi F 2000 Round-off errors and *p*-adic numbers *Nonlinearity* **13** 309–22
- [11] Bruin H, Lambert A, Poggiaspalla G and Vaienti S 2003 Numerical analysis for a discontinuous rotation of the torus Chaos 13 558–71
- [12] Buzzi J 2001 Piecewise isometries have zero topological entropy Ergodic Theory Dyn. Syst. 21 1371-7
- [13] DeVogelaere R 1958 On the structure of symmetric periodic solutions of conservative systems, with applications Contributions to The Theory of Nonlinear Oscillations vol 4 (Annals of Mathematics Studies vol 41) (Princeton, NJ: Princeton University Press) pp 53–84
- [14] Goetz A 2002 Piecewise isometries—An emerging area of dynamical systems Fractals in Graz 2001 ed P Grabner and W Woess (Basel: Birkhäuser) pp 135–44
- [15] Goetz A and Poggiaspalla G 2004 Rotations by $\pi/7$ Nonlinearity 17 1787–802
- [16] Huxley M N 1988 The area within a curve Proc. Indian Acad. Sci. Math. Sci. 97 111-6
- [17] Koupstov K, Lowenstein J H and Vivaldi F 2002 Quadratic rational rotations of the torus and dual lattice maps Nonlinearity 15 1795–842
- [18] Kozyakin V, Kuznetsov N, Pokrovskii A and Vladimirov I 1997 Some problems in analysis of discretizations of continuous dynamical systems Proc. 2nd World Congress of Nonlinear Analysts, Part 2 (Athens, Greece, 1996) vol 30, pp 767–78
- [19] Lowenstein J H, Koupstov K L and Vivaldi F 2004 Recursive tiling and geometry of piecewise rotations by π/7 Nonlinearity 17 371–95
- [20] Lowenstein J H, Hatjispyros S and Vivaldi F 1997 Quasi-periodicity, global stability and scaling in a model of Hamiltonian round-off Chaos 7 49–56
- [21] Surer P 2009 ϵ -shift radix systems and radix representations with shifted digit sets *Publ. Math. Debrecen* 74 19–43
- [22] Vinogradov I M 1954 Elements of Number Theory (New York: Dover)
- [23] Vinogradov I M 1985 Selected Works (Berlin: Springer)
- [24] Vivaldi F 1994 Periodicity and transport from round-off errors Exp. Math. 3 303-15
- [25] Vivaldi F 2006 The arithmetic of discretized rotations, in p-adic Mathematical Physics AIP Conf. Proc. 826 162–73
- [26] Vivaldi F and Vladimirov I 2003 Pseudo-randomness of round-off errors in discretized linear maps on the plane Int. J. Bifurcations Chaos 13 3373–93
- [27] Vladimirov I 1996 Quantized linear systems on integer lattice: frequency-based approach. Part 1 CADSEM Report 96-032 (Melbourne: Deakin University) pp 1–37
 - Vladimirov I 1996 Quantized linear systems on integer lattice: frequency-based approach. Part 2 *CADSEM Report* 96-033 (Melbourne: Deakin University) pp 1–50