Problems by Attila Pethő, Debrecen

In this talk I collected some problem, which I proposed and/or tried to solve during my career.

1. Powers in linear recursive sequences

To find perfect powers and polynomial values in lrs is one of my favorite topics. A long standing problem was to prove that 0, 1, 8 and 144 are the only powers in the Fibonacci sequence. This was proved finally by Bugeaud, Mignotte and Siksek in 2006.

In 1996 at The Seventh International Research Conference on Fibonacci Numbers and Their Applications I proposed the following [16]

Problem 1.: The sequence of tribonacci numbers is defined by $T_0 = T_1 = 0, T_2 = 1$ and $T_{n+3} = n + 2 + T_{n+1} + T_n$ for $n \ge 0$. Are the only squares $T_0 = T_1 = 0, T_2 = T_3 = 1, T_5 = 4, T_{10} = 81, T_{16} = 3136 = 56^2$ and $T_{18} = 10609 = 103^2$ in T_n ?

By using the sieving moduli 3, 7, 11, 13, 29, 41, 43, 53, 79, 101, 103, 131, 239, 97, 421, 911, 1021 and 1123 one can show that this is true for $n \leq 2 \cdot 10^6$, but known methods do not seems to be applicable for the solution of this problem.

The problem is still unsolved, although in the edited version of the second part of that talk [17] combining results of Shorey and Stewart [22] with that of Corvaja and Zannier [11] I proved

Theorem 1. Let G_n be a third order LRS. For the roots α_i , i = 1, 2, 3 of the characteristic polynomial of G_n assume that $|\alpha_1| > |\alpha_2| \ge |\alpha_3|$ and non of them is a root of unity. Then there are only finitely many perfect powers in G_n .

As the characteristic polynomial of the tribonacci sequence $x^3 - x^2 - x - 1$ is irreducible with one dominating real root ≈ 1.839286755 it follows that there exist finitely many perfect powers in it. Unfortunately the proof of the theorem is only partially effective, we have an effective bound for the exponent of the possible perfect powers, but no effective bound for the size of a fixed power, e.g. for squares.

I think that Theorem 1 can be generalized at least in the following form:

Problem 2.: Let G_n be an LRS such that its characteristic polynomial is irreducible and has a dominating root, then there is only finitely many perfect powers in it.

By a result of Shorey and Stewart [22] the exponent of perfect powers can be bounded effectively, the problem is to handle the powers with bounded exponent. Combining this with the result of Corvaja and Zannier [11] and with the combinatorics of the roots, like in Pethő [17], one can probably setle this conjecture.

Like the Fibonacci sequence, we can continue the tribonacci sequence in "negative direction", and get $T_{-n} = -T_{-n+1} - T_{-n+2} + T_{-n+3}$ with initial terms $T_0 = 0, T_{-1} = 1, T_{-2} = -1$. We call this sequence n-tribonacci. One can again ask, which are the perfect powers in this sequence. After a simple search we find: $T_0 = T_{-3} = T_{-16} = 0, T_{-1} = T_{-6} = -T_{-2} = 1, T_{-7} = 2^2, T_{-8} = (-2)^3, T_{-13} = 3^2, T_{-29} = 3^4, T_{-32} = 56^2, T_{-33} = 103^2$ and $T_{-62} = 6815^2$. It is interesting to observe that $T_{10} = T_{-29}, T_{16} = T_{-32}$ and $T_{18} = T_{-33}$.

Problem 3.: Are all perfect powers of the n-tribonacci sequence listed above? Are there only finitely many perfect powers in the n-tribonacci sequence?

The answer seems to be very difficult, because the characteristic polynomial of the n-tribonacci sequence has two conjugate complex roots of the same absolute value and its real root is less than one.

Let $a, b \in \mathbb{Z}$ and $\delta \in \{1, -1\}$ such that $a^2 - 4(b - 2\delta) \neq 0$, $b\delta \neq 2$ and if $\delta = 1$ then $b \neq 2a - 2$. Let further the sequence $G_n = G_n(a, b, \delta)$, $n \geq 0$ defined by the initial terms $G_0 = 0$, $G_1 = 1$, $G_2 = a$, $G_3 = a^2 - b - \delta$ and by the recursion

(1)
$$G_{n+4} = aG_{n+3} - bG_{n+2} + \delta aG_{n+1} - G_n, \quad n \ge 0.$$

I proved in [18] that these are divisibility sequences, i.e., $G_n|G_m$, whenever n|m. More precisely, the roots of the characteristic polynomial of G_n can be numbered so that they are $\eta, \frac{\delta}{\eta}, \vartheta, \frac{\delta}{\vartheta}$ and

$$G_n = \frac{\eta^n - \vartheta^n}{\eta - \vartheta} \frac{1 - \left(\frac{\delta}{\eta\vartheta}\right)^n}{1 - \frac{\delta}{\eta\vartheta}}$$

Here we ask again to prove that for fixed a, b there are only finitely many perfect powers in G_n . We can again bound the exponent, but can not treat the equation $G_n = x^q$ for fixed q > 1. Especially complicated seems the case q = 2, because the gcd of the divisors $\frac{\eta^n - \vartheta^n}{\eta - \vartheta}$ and $\frac{1 - \left(\frac{\delta}{\eta \vartheta}\right)^n}{1 - \frac{\delta}{\eta \vartheta}}$ can be arbitrary large.

2. Thue equations

After the work of E. Thomas [23] several paper appeared about the solutions of parametrized families of Thue equations. With Halter-Koch, Lettl and Tichy we proved [13] the following: **Theorem 2.** Let $n \geq 3$, $a_1 = 0, a_2, \ldots, a_{n-1}$ be distinct integers and $a_n = a$ an integral parameter. Let $\alpha = \alpha(a)$ be a zero of $P(x) = \prod_{i=1}^{n} (x - a_i) - d$ with $d = \pm 1$ and suppose that the index I of $\langle \alpha - a_1, \ldots, \alpha - a_{n-1} \rangle$ in $U_{\mathcal{O}}$, the group of units of \mathcal{O} , is bounded by a constant $J = J(a_1, \ldots, a_{n-1}, n)$ for every a from some subset $\Omega \subset \mathbb{Z}$. Assume further that the Lang-Waldschmidt conjecture is true. Then for all but finitely many values $a \in \Omega$ the diophantine equation

(2)
$$\prod_{i=1}^{n} (x - a_i y) - dy^n = \pm 1$$

only has trivial solutions, except when n = 3 and $|a_2| = 1$, or when n = 4and $(a_2, a_3) \in \{(1, -1), (\pm 1, \pm 2)\}$, in which cases (2) has exactly one more general solution.

The assumption on the index I is technical, the essential assumption is the Land-Waldschmidt conjecture. In the cited paper we formulated:

Problem 4.: The last theorem is true for all large enough parameter value without further assumptions.

A weaker version of this conjecture was formulated by E. Thomas [24]. He assumed that $a_i = p_i(a), i = 2, ..., n-1$ and $0 < \deg p_2 < \cdots < \deg p_{n-1}$, where p_i denotes monic polynomial with integer coefficients. This weaker conjecture was proved by C. Heuberger [14] under some technical conditions on the degree of the polynomials.

3. Progressions in the set of solutions of norm form equations

Let \mathbb{K} be an algebraic number field of degree k, and let $\alpha_1, \ldots, \alpha_n$ be linearly independent elements of $\mathbb{Z}_{\mathbb{K}}$ over \mathbb{Q} . Let m be a non-zero integer and consider the norm form equation

(3)
$$N_{\mathbb{K}/\mathbb{Q}}(x_1\alpha_1 + \ldots + x_n\alpha_n) = m$$

in integers x_1, \ldots, x_n . Let H denote the solution set of (3) and |H| the size of H. Note that if the \mathbb{Z} -module generated by $\alpha_1, \ldots, \alpha_n$ contains a submodule, which is a full module in a subfield of $\mathbb{Q}(\alpha_1, \ldots, \alpha_n)$ different from the imaginary quadratic fields and \mathbb{Q} , then this equation can have infinitely many solutions (see e.g. Schmidt [21]).

Arranging the elements of H in an $|H| \times n$ array \mathcal{H} , one may ask at least two natural questions about arithmetical progressions appearing in H. The "horizontal" one: do there exist infinitely many rows of \mathcal{H} , which form arithmetic progressions; and the "vertical" one: do there exist arbitrary long arithmetic progressions in some column of \mathcal{H} ? Note that the first question is meaningful only if n > 2.

We are now presenting an example. Let $\mathbb{K} := \mathbb{Q}(\alpha)$ with $\alpha^5 = 3$. Then

$$N_{\mathbb{K}/\mathbb{Q}}(x_1 + x_2\alpha + \dots + x_5\alpha^4) = 9x_3^5 + 81x_5^5 + x_1^5 + 27x_4^5 + 3x_2^5 - 135x_5^3x_4x_1 + 45x_5x_2x_4^2x_1^2 + 135x_2x_4^2x_2^2 - 45x_2x_4^3x_1 + 45x_5x_3x_1^2 - 45x_2x_3^3x_4 + 135x_3^2x_5^2x_4 + 45x_1x_5^2x_2^2 - 45x_4x_2^3x_5 + 45x_4^2x_2^2x_3 + 45x_4^2x_1x_3^2 - 15x_4x_1^3x_3 + 15x_4x_1^2x_2^2 + 15x_2x_3^2x_1^2 + 45x_5x_2^2x_3^2 - 15x_5x_1^3x_2 - 135x_5x_3x_4^3 - 135x_2x_5^3x_3 - 45x_5x_3^3x_1 - 15x_2^3x_3x_1 - 45x_2x_5x_3x_4x_1.$$

The next table contains some solution of the equation $N_{\mathbb{K}/\mathbb{Q}}(x_1 + x_2\alpha + \dots + x_5\alpha^4) = 1$:

x_1	x_2	x_3	x_4	x_5
4	-5	4	-2	0
1	2	-1	-1	0
4	2	0	0	1
1	1	0	1	0
1	5	1	2	2
-17	1	-6	3	8
7	6	5	4	3
-2	-1	1	1	0
-11	-5	5	6	0
-2	0	1	-1	1
-8	-8	1	6	2
28	16	4	3	8
10	12	12	4	9
•••	• • •			

The bold numbers form a five term horizontal AP and a seven terms vertical AP. The "horizontal" problem was treated by Bérczes and Pethő [8] by proving that if $\alpha_i = \alpha^{i-1}$ (i = 1, ..., n) then in general \mathcal{H} contains only finitely many effectively computable "horizontal" AP's and they were able to localize the possible exceptional cases. The following question remains unanswered:

Problem 5.: Does there exist infinitely many quartic integers α such that $\frac{4\alpha^4}{\alpha^4-1} - \frac{\alpha}{\alpha-1}$ is a quadratic algebraic number.

We were able to found only one example with defining polynomial $x^4 + 2x^3 + 5x^2 + 4x + 2$ such that the corresponding element is a real quadratic number. It is a root of $x^2 - 4x + 2$. Allowing however α not to be integral we can obtain a lot of examples.

It seems to be a simple problem to extend the result of Bérczes and Pethő to geometric progressions. This is equivalent to investigate the equation

$$N_{\mathbb{K}/\mathbb{Q}}\left(\frac{\alpha^n q^n - 1}{\alpha q - 1}\right) = m,$$

where $\mathbb{K} = \mathbb{Q}(\alpha)$ and α denotes an algebraic integer of degree *n*. I expect that if n > 1 then this equation has only finitely many solutions in $q \in \mathbb{Z}$.

The investigation of the "vertical" AP's is much more difficult. In this direction Bérczes, Hajdu and Pethő [9]proved

Theorem 3. Let $(x_1^{(j)}, \ldots, x_n^{(j)})$ $(j = 1, \ldots, t)$ be a sequence of distinct elements in H such that $x_i^{(j)}$ is a non-zero arithmetic progression for some $i \in \{1, \ldots, n\}$. Then we have $t \leq c_1$, where $c_1 = c_1(k, m)$ is an explicitly computable constant.

It is interesting to note that c_1 is independent from the field. One can probably strength this result such that the upper bound for the length of the AP's depend not on m, but only on the number of its prime divisors. It is even possible that the bound depends only on k.

Earlier Pethő and Ziegler [20] as well as Dujella, Pethő and Tadić [10] investigated the AP's on Pell equations. They proved that for all but one non-constant AP of integers of length four y_1, y_2, y_3, y_4 there exist infinitely many integers d, m for which $x_i^2 - dy_i^2 = m, i = 1, 2, 3, 4$ with some integers $x_i = x_i(d, m, y_1, \ldots, y_4), i = 1, 2, 3, 4$. In contrast, five term AP's are lying on only finitely many Pell equations.

Problem 6.: Prove analogous result for norm form equations over cubic number fields. More specifically: let $y^{(i)}$, i = 1, ..., 5 an AP of integers. Then there exist infinitely many $m \in \mathbb{Z}$ and \mathbb{Q} -independent algebraic integers $\alpha_1, \alpha_2, \alpha_3$ such that $\mathbb{K} = \mathbb{Q}(\alpha_2, \alpha_3)$ has degree three and (3) holds for $(x_1^{(i)}, x_2^{(i)}, y^{(i)}), i = 1, ..., 5$ with some $x_1^{(i)}, x_2^{(i)}\mathbb{Z}$. Can 5 replace with a larger number?

In the above mentioned papers we worked out a systematic method to find Pell equations having long AP's. For example the AP -7, -5, -3, -1, 1, 3, 5, 7 is lying on the equation $x^2 - 570570y^2 = 4406791$ and -461, -295, -129, 37, 203, 369, 535 on $x^2 + 1245y^2 = 375701326$.

Problem 7.: Find a systematic method to construct cubic norm form equations with long AP. Do the same for higher degree norm form equations.

Problem 8.: Prove analogous results for geometric progressions.

4. Polynomials

Problem 9.: Let K be a algebraically closed field of characteristic zero. Characterize all $P(X) \in K[X], Q(Y) \in K[Y], R(X, Y) \in K[X, Y]$ such that the set of zeroes of P(X) and Q(Y) coincide, provided R(X, Y) = 0.

The case R(X,Y) = Y - A(X) was solved completely by Fuchs, Pethő and Tichy [12]. They proved

Theorem 4. Assume that P(X) has k different zeroes. Then there exist $a, b, c \in K, a, c \neq 0$ such that: if k = 1 then

$$P(X) = a(x-b)^{\deg P} \text{ and } A(X) = c(x-b)^{\deg A} + b;$$

if $k \geq 2$ then either A(X) = X or $A(X) = ax + b, a \neq 1$ and in this case

$$P(X) = c\left(X + \frac{b}{a-1}\right)^{s} \prod_{i=1}^{r} \prod_{j=0}^{\ell-1} \left(X - a^{j}x_{i} - b\frac{a^{j}-1}{a-1}\right),$$

where x_1, \ldots, x_r are all different and ℓ is the multiplicative order of a.

5. Shift radix systems

For $(r_1, \ldots, r_d) = \mathbf{r} \in \mathbb{R}^d$ and $\mathbf{a} = (a_1, \ldots, a_d) \in \mathbb{Z}^d$ let $\tau_{\mathbf{r}}(\mathbf{a}) = (a_2, \ldots, a_d, -\lfloor \mathbf{ra} \rfloor)^T$, where \mathbf{ra} denotes the scalar product. This nearly linear mapping was introduced by Akiyama, Borbély, Brunotte, Thuswaldner and myself [1]. We proved that it can be considered as a common generalization of canonical number systems (CNS) and β -expansions.

We also defined the sets

 $\mathcal{D}_d = \{ \mathbf{r} : \{ \tau_{\mathbf{r}}^k(\mathbf{a}) \}_{k=0}^{\infty} \text{ is bounded for all } \mathbf{a} \in \mathbb{Z}^d \}, \\ \mathcal{D}_d^0 = \{ \mathbf{r} : \{ \tau_{\mathbf{r}}^k(\mathbf{a}) \}_{k=0}^{\infty} \text{ is ultimately zero for all } \mathbf{a} \in \mathbb{Z}^d \}$

and \mathcal{E}_d , which is the set of real monic polynomials, whose roots are lying in the closed unit disc. We proved in the same paper that if $\mathbf{r} \in \mathcal{D}_d$ then $R(X) = X^d + r_d X^{d-1} + \cdots + r_2 X + r_1 \in \mathcal{E}_d$ and if R(X) is lying in the interior of \mathcal{E}_d then $\mathbf{r} \in \mathcal{D}_d$.

We called $\tau_{\mathbf{r}}$ a shift radix system (SRS), if $\mathbf{r} \in \mathcal{D}_d^0$ and gave an algorithm, which decides whether $\mathbf{r} \in \mathbb{Q}^d$ is a SRS. However this algorithm is exponential, moreover we are not able to give a polynomial time verification for $\mathbf{r} \notin \mathcal{D}_d^0 \cap \mathbb{Q}^d$. We found points $\mathbf{r} \in \mathbb{Q}^2$ such that $\mathbf{r} \notin \mathcal{D}_2^0$ and the cycles proving this can be arbitrary long. Computational experiments, see e.g. [1, 15] support the following :

Problem 10.: Prove that the SRS problem can not be solved by a polynomial time algorithm. Stronger statement is that it does not belong to the NP complexity class.

The structure of \mathcal{D}_d^0 , especially approaching its boundary is very complicated, see [2] for d = 2. On the other hand we know [1], that the closure of \mathcal{D}_d is \mathcal{E}_d . However the investigation of the boundary points of \mathcal{E}_d leads to interesting and hard problems. The case d = 2 was studied by Akiyama et al in [2]. They proved that \mathcal{D}_2 is equal to the closed triangle with vertices (-1,0), (1,-2), (1,2), but without the points (1,-2), (1,2), the line segment $\{(x, -x - 1) : 0 < x < 1\}$ and, possibly, some points of the line segment $\{(1,\lambda) : -2 < \lambda < 2\}$. Write in the last case $\lambda = 2\cos\alpha$ and $\omega = \cos\alpha + i\sin\alpha$. It is easy to see, that if $\lambda = 0, \pm 1$ (i.e. $\alpha = 0, \pm \pi/2$) then $(1, \lambda)$ belongs to \mathcal{D}_2 and we conjectured in [2] that this is true for all points of the line segment. In [5] the conjecture was proved for the golden mean, i.e. for $\lambda = \frac{1+\sqrt{5}}{2}$ and in [6] for those ω , which are quadratic algebraic numbers. The conjecture has the following nice arithmetical form:

Problem 11.: Let $|\lambda| < 2$ be a real number. If the sequence of integers $\{a_n\}$ satisfies the relation

$$0 \le a_{n-1} + \lambda a_n + a_{n+1} < 1$$

then it is periodic.

If ω , defined above is a root of unity then the problem may be easier as in the general case. On the other hand from the point of view of arithmetic the cases, when λ is a rational number, e.g., $\lambda = \frac{1}{2}$ seems simpler.

If the point \mathbf{r} is lying on the boundary of \mathcal{E}_d then either $\mathbf{r} \in \mathcal{D}_d$ or $\mathbf{r} \notin \mathcal{D}_d$. With other words this means that the sequence $\{\tau_{\mathbf{r}}(\mathbf{a})\}$ is ultimately periodic for all $\mathbf{a} \in \mathbb{Z}^d$ and there exists $\mathbf{a} \in \mathbb{Z}^d$ for which it is divergent. Presently we do not know any general method to distinguish between these cases. Recently I gave an algorithm [19] in the special case, when $\pm 1, \pm i$ is a simple root of $X^d + r_d X^{d-1} + \cdots + r_2 X + r_1$.

Problem 12.: Is it algorithmically decidable for $\mathbf{r} \in \mathcal{E}_d$ whether $\mathbf{r} \in \mathcal{D}_d$?

I am not sure that the answer is affirmative. The problem is open even for d = 2. In this case, by the results of [2], the status only points of the line segment $\{(1, y) : -2 < y < 2\}$ is questionable. If the answer to Problem 9 is affirmative, which I strongly believe, then d = 2 would be completely solved. A related, probably easier problem is:

Problem 13.: Prove that there are lying no elements of \mathcal{D}_d^0 on the boundary of \mathcal{E}_d .

This is true for d = 2 [2], but open for $d \ge 3$.

For each $d \in \mathbb{N}, d \ge 1$ define the set

 $\mathcal{B}_d = \{(b_1, \dots, b_d) \in \mathbb{Z}^d : X^d - b_1 X^{d-1} - \dots - b_d \text{ is a Pisot or Salem polynomial}\}.$ Further for $M \in \mathbb{N}_{>0}$ set

(4)
$$\mathcal{B}_d(M) = \left\{ (b_2, \dots, b_d) \in \mathbb{Z}^{d-1} : (M, b_2, \dots, b_d) \in \mathcal{B}_d \right\}.$$

It is clear that $\mathcal{B}_d(M)$ is a finite set. In [3] we proved

Theorem 5. Let $d \ge 2$. We have

(5)
$$\left| \frac{|\mathcal{B}_d(M)|}{M^{d-1}} - \lambda_{d-1}(\mathcal{D}_{d-1}) \right| = O(M^{-1/(d-1)}),$$

where λ_{d-1} denotes the (d-1)-dimensional Lebesgue measure.

$$\hat{\mathcal{B}}_d(M) = \left\{ (b_1, b_2, \dots, b_d) \in \mathbb{Z}^d \cap \mathcal{B}_d : \max\{|b_1|, |b_2|, \dots, |b_d|\} \le M \right\}.$$

Problem 14.: Does there exist a constant c, such that

$$\lim_{M \to \infty} \frac{|\hat{\mathcal{B}}_d(M)|}{M^d} = c?$$

References

- S. AKIYAMA, T. BORBÉLY, H. BRUNOTTE, A. PETHŐ AND J. M. THUSWALDNER, Generalized radix representations and dynamical systems I, Acta Math. Hungar., 108 (2005), 207–238.
- [2] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ AND J. M. THUSWALDNER, Generalized radix representations and dynamical systems II, Acta Arith. 121 (2006), 21-61.
- [3] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ AND J. M. THUSWALDNER, Generalized radix representations and dynamical systems IV, Indag. math. to appear.
- [4] H. BRUNOTTE, On trinomial bases of radix representations of algebraic integers, Acta Sci. Math. (Szeged), 67 (2001), 521-527.
- [5] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ and W. Steiner, Remarks on a conjecture on certain integer sequences, Periodica Mathematica Hungarica, 52 (2006), 1-17.
- [6] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ and W. STEINER, Periodicity of certain piecewise affine planar maps, Tsukuba J. Math., 32 (2008), 197-251.
- [7] Y. BUGEAUD, M. MIGNOTTE and S. SIKSEK, Classical and modular approaches to exponential Diophantine equations. I. Fibonacci and Lucas perfect powers., Annals of Math. 163 (2006), 969-1018.
- [8] A. BÉRCZES and A. PETHŐ, On norm form equations with solutions forming arithmetic progressions, Publ. Math. Debrecen, 65 (2004), 281-290.
- [9] A. BÉRCZES, L. HAJDU and A. PETHŐ, Arithmetic progressions in the solution sets of norm form equations, Rocky Mountain J. Math., to appear.
- [10] A. DUJELLA, A. PETHŐ and P. TADIĆ, On arithmetic progressions on Pellian equations, Acta Math. Hungar., 120 (2008), 29-38.
- [11] P. CORVAJA and U. ZANNIER, Some new applications of the subspace theorem., Compositio Math. 131 (2002), 319–340.
- [12] CL. FUCHS, A. PETHŐ and R.F. TICHY, On the diophantine equation $G_n(x) = G_m(P(x))$: higher order linear recurrences, Trans. Amer. Math. Soc., **355** (2003), 4657–4681.
- [13] F. HALTER-KOCH, G. LETTL, A. PETHŐ and R.F. TICHY, Thue Equations associated with Ankeny-Brauer-Chowla Number Fields, J. London Math. Soc. (2) 60 (1999), 1– 20.
- [14] C. HEUBERGER, On a conjecture of E. Thomas concerning parametrized Thue equations., Acta Arith. 98 (2001), 375394.
- [15] A. KOVÁCS, Generalized binary number systems, Ann. Univ. Sci. Budap. Rolando Eötvös, Sect. Comput., 20 (2001), 195-206.
- [16] A. PETHŐ, Diophantine properties of linear recurrence sequences I. In: "Applications of Fibonacci Numbers, Volume 7", Ed.: G.E. Bergum, Kluwer Academic Publishers, 1998, pp. 295–309.
- [17] A. PETHŐ, Diophantine properties of linear recursive sequences. II., Acta. Math. Acad. Paed. Nyíregyháziensis, 17 (2001), 81-96.
- [18] A. PETHŐ, Egy negyedrendű rekurzív sorozatcsaládról, Acta Acad. Paed. Agriensis, Sect. Math., 30 (2003), 115-122.
- [19] A. PETHŐ, On the boundary of the set of the closure of contractive polynomials, Integers, to appear.
- [20] A. PETHŐ and V. ZIEGLER, Arithmetic progressions on Pell equations, J. Number Theory, 128 (2008), 1389-1409.

- [21] W.M. SCHMIDT, Norm form equations, Ann. of Math., 96 (1972), 526–551.
- [22] T.N. SHOREY and C.L. STEWART, On the Diophantine equation $ax^{2t} + bx^ty + cy^2 = d$ and pure powers in recurrences, Math. Scand., **52** (1983), 24–36.
- [23] E. THOMAS, Complete solutions to a family of cubic Diophantine equations. J. Number Theory 34 (1990), 235–250.
- [24] E. THOMAS, Solutions to certain families of Thue equations, J. Number Theory 43 (1993), 319369.